


# ASCENT-04: Analysis of Efficacy by Biomarker Subgroups With Sacituzumab Govitecan + Pembrolizumab vs Chemotherapy + Pembrolizumab in Participants With Previously Untreated PD-L1+ Metastatic Triple-Negative Breast Cancer

Sara M Tolaney<sup>1</sup>, Peter Schmid<sup>2</sup>, Evandro de Azambuja<sup>3</sup>, Kevin Kalinsky<sup>4</sup>, Sung-Bae Kim<sup>5</sup>, Clinton Yam<sup>6</sup>, Bernardo Rapoport<sup>7,8</sup>, Seock-Ah Im<sup>9</sup>, Barbara Pistilli<sup>10</sup>, Wassim Mchayleh<sup>11</sup>, David W Cescon<sup>12</sup>, Junichiro Watanabe<sup>13</sup>, Manuel Alejandro Lara Bañuelas<sup>14</sup>, Ruffo Freitas-Junior<sup>15</sup>, Alain Lortholary<sup>16</sup>, Catherine Lai<sup>17</sup>, Ann Chen<sup>17</sup>, Meghna Das Thakur<sup>17</sup>, Yajia Zhang<sup>17</sup>, Sherene Loi<sup>18</sup>

<sup>1</sup>Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA, USA; <sup>2</sup>Centre for Experimental Cancer Medicine, Bart's Cancer Institute, Queen Mary University of London, London, UK; <sup>3</sup>Institut Jules Bordet, Hôpital Universitaire de Bruxelles (H.U.B), Université Libre de Bruxelles (U.L.B.), Brussels, Belgium; <sup>4</sup>Winship Cancer Institute, Emory University, Atlanta, GA, USA; <sup>5</sup>Asan Medical Center, University of Ulsan College of Medicine, Seoul, Republic of Korea; <sup>6</sup>The University of Texas MD Anderson Cancer Center, Houston, TX, USA; <sup>7</sup>The Medical Oncology Centre of Rosebank, Clinical and Translational Research Unit (CTRU), Saxonworld, South Africa; <sup>8</sup>Department of Immunology, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa; <sup>9</sup>Seoul National University Hospital, Cancer Research Institute, Seoul National University College of Medicine, Seoul National University, Seoul, Republic of Korea; <sup>10</sup>Gustave Roussy, IHU-National PReclSion Medicine Center in Oncology, Villejuif, France; <sup>11</sup>AdventHealth Cancer Institute, Orlando, FL, USA; <sup>12</sup>Princess Margaret Cancer Centre/UHN, Toronto, ON, Canada; <sup>13</sup>Juntendo University Graduate School of Medicine, Tokyo, Japan; <sup>14</sup>SCIENTIA Investigación Clínica, Chihuahua, Mexico; <sup>15</sup>Advanced Center for Diagnosis of Breast Diseases Federal University of Goiás, Goiás, Brazil; <sup>16</sup>Groupe d'Investigateurs National des Etudes des Cancers Ovariens et du sein (GINECO) and Hôpital Privé du Confluent, Nantes, France; <sup>17</sup>Gilead Sciences, Inc., Foster City, CA, USA; <sup>18</sup>Peter MacCallum Cancer Centre, Melbourne, Australia

# Introduction

- The ASCENT-04 study demonstrated significant, clinically meaningful improvement in PFS with SG + pembro vs chemo + pembro (HR, 0.65; 95% CI, 0.51-0.84;  $P < .001$ ) in participants with previously untreated PD-L1–positive advanced TNBC<sup>1</sup>
- SG targets Trop-2, a surface protein overexpressed in many solid tumors, particularly in TNBC<sup>2-4</sup>
- Mutations in *BRCA1* and *BRCA2* and HER2 amplification or overexpression are associated with altered efficacy of some mBC treatments, although the role of these biomarkers is less established in PD-L1–positive TNBC<sup>5-9</sup>



Prespecified retrospective exploratory analyses evaluated the impact of Trop-2, BRCA, and HER2 biomarkers on efficacy of SG + pembro vs chemo + pembro in ASCENT-04

**Chemo**, chemotherapy; **HER2**, human epidermal growth factor receptor 2; **HR**, hazard ratio; **mBC**, metastatic breast cancer; **PD-L1**, programmed cell death ligand 1; **pembro**, pembrolizumab; **PFS**, progression-free survival; **SG**, sacituzumab govitecan; **TNBC**, triple-negative breast cancer.

1. Tolaney SM, et al. *N Engl J Med*. 2026;394:354-66. 2. Goldenberg DM, et al. *Oncotarget*. 2015;6:22496-512. 3. Jacot W, et al. *Cancer Med*. 2025;14:e70615. 4. Zhao W, et al. *Oncol Rep*. 2018;40:759-66. 5. Godet I, et al. *Integr Cancer Sci Ther*. 2017;4:10.15761/ICST.1000228. 6. Stoppa-Lyonnet D. *Eur J Hum Genet*. 2016;24:S3-S9. 7. Bardia A, et al. *Ann Oncol*. 2021;32:1148-56. 8. McCann KE, et al. *Drugs Context*. 2018;7:212540. 9. Chen W, et al. *Cancer Discov*. 2026;16:235-49.

# Methods

- Participants with previously untreated, locally advanced unresectable or metastatic PD-L1–positive (PD-L1 CPS<sup>a</sup> ≥ 10) TNBC were randomized 1:1 to receive SG + pembro or chemo + pembro; the primary end point was PFS by BICR
- This prespecified exploratory analysis was performed using central testing of fresh or archival (formalin-fixed paraffin-embedded) tumor samples; 48% from metastatic sites in the Trop-2 biomarker analysis set
- Biomarker status was analyzed descriptively for association with PFS by BICR
- At the primary data cutoff (March 2025), median follow-up was 14.0 months (range, 0.1-28.6)<sup>1</sup>
- Participants from the ITT population were included in the biomarker analysis set if they had ≥ 1 evaluable biomarker measurement available at baseline

## Subgroups Analyzed

### Trop-2 Expression

- Characterized by H-score, as determined by IHC<sup>b</sup>
- Participants grouped by Trop-2 expression quartile

### tBRCA Status


- Measured by WES as local testing was uncommon
- Participants grouped by tBRCA WT or mut status
- Mut indicates mutation in *BRCA1*, *BRCA2*, or both genes

### HER2 Expression

- Measured by ISH and IHC
- Participants grouped as HER2 IHC 0 or HER2 low
- HER2 low includes IHC 1+ or IHC 2+/ISH–

<sup>a</sup>Measured per the 22C3 assay. <sup>b</sup>Scoring as follows: 0, negative or no staining of tumor cell; 1, weak or faint staining; 2, moderate staining; 3, strong staining. H-score determined by (% of 1+ tumor cells) + (2x% of 2+ tumor cells) + (3x% of 3+ tumor cells). BICR, blinded independent central review; chemo, chemotherapy; CPS, combined positive score; HER2, human epidermal growth factor receptor 2; IHC, immunohistochemistry; ITT, intent-to-treat; mut, mutation; PD-L1, programmed cell death ligand 1; pembro, pembrolizumab; PFS, progression-free survival; SG, sacituzumab govitecan; TNBC, triple-negative breast cancer; WT, wild-type.  
1. Tolaney SM, et al. *N Engl J Med*. 2026;394:354-66.

# PFS by Trop-2 H-score Quartiles<sup>a</sup>







PFS was longer with SG + pembro vs chemo + pembro across all Trop-2 subgroups

<sup>a</sup>Trop-2 H-score quartiles: Q1, 0-224; Q2, 225-279; Q3, 280-298; Q4, 299-300. <sup>b</sup>HR value is stratified. <sup>c</sup>HR values are unstratified.

BAS, biomarker analysis set; chemo, chemotherapy; HR, hazard ratio; ITT, intent-to-treat; mo, months; mPFS, metastatic progression-free survival; NE, not estimable; pembro, pembrolizumab; PFS, progression-free survival; Q, quartile; SG, sacituzumab govitecan.

1. Tolaney SM, et al. *N Engl J Med.* 2026;394:354-66.

# PFS by Trop-2 H-Score Quartiles<sup>a</sup>




SG + pembro treatment was associated with consistently improved PFS vs chemo + pembro across Trop-2 expression quartiles, with trends toward greater separation of KM curves in Q3 and Q4

<sup>a</sup>Trop-2 H-score quartiles: Q1, 0-224; Q2, 225-279; Q3, 280-298; Q4, 299-300.

BICR, blinded independent central review; chemo, chemotherapy; HR, hazard ratio; KM, Kaplan-Meier; mo, months; NE, not estimable; pembro, pembrolizumab; PFS, progression-free survival; Q, quartile; SG, sacituzumab govitecan.

# PFS by tBRCA Subgroups



- From the biomarker analysis set, 23% of participants in the SG + pembro group and 20% in the chemo + pembro group had tBRCA mut tumors


Median PFS was longer with SG + pembro vs chemo + pembro in both tBRCA subgroups

<sup>a</sup>HR value is stratified. <sup>b</sup>HR values are unstratified.


BAS, biomarker analysis set; chemo, chemotherapy; 2; HR, hazard ratio; ITT, intent-to-treat; mo, months; mPFS, median progression-free survival; mut, mutant; NE, estimable; pembro, pembrolizumab; PFS, progression-free survival; SG, sacituzumab govitecan; tBRCA, tumor BRCA; TNBC, triple-negative breast cancer; WES, whole exome sequencing; WT, wild-type.

1. Tolaney SM, et al. *N Engl J Med.* 2026;394:354-66.

# PFS by tBRCA Subgroups



No. at risk	0	2	4	6	8	10	12	14	16	18	20	22	24	26
WT, SG + pembro	130	119	100	78	58	42	30	19	17	11	7	3	0	0
WT, Chemo + pembro	131	113	92	71	47	30	21	15	13	8	4	2	1	0




No. at risk	0	2	4	6	8	10	12	14	16	18	20	22	24	26
Mut, SG + pembro	39	36	33	28	21	16	11	10	8	6	2	1	1	0
Mut, Chemo + pembro	32	30	29	22	19	14	9	6	2	0	0	0	0	0

PFS improvement with SG + pembro vs chemo + pembro was observed across tBRCA subgroups

BICR, blinded independent central review; chemo, chemotherapy; HR, hazard ratio; mBC, metastatic breast cancer; mo, months; mut, mutant; NE, not estimable; pembro, pembrolizumab; PFS, progression-free survival; SG, sacituzumab govitecan; tBRCA, tumor BRCA; WT, wild-type.

# PFS by HER2 Subgroups




Median PFS was longer with SG + pembro vs chemo + pembro across both HER2 subgroups

<sup>a</sup>HR value is stratified. <sup>b</sup>HR values are unstratified.


**BAS**, biomarker analysis set; **chemo**, chemotherapy; **HER2**, human epidermal growth factor receptor 2; **HR**, hazard ratio; **IHC**, immunohistochemistry; **ITT**, intent-to-treat; **mo**, months; **mPFS**, median progression-free survival; **pembro**, pembrolizumab; **PFS**, progression-free survival; **SG**, sacituzumab govitecan.

1. Tolaney SM, et al. *N Engl J Med*. 2026;394:354-66.

# PFS by HER2 Subgroups



No. at risk	0	2	4	6	8	10	12	14	16	18	20	22	24	26
HER2 IHC 0, SG + pembro	84	76	64	54	42	32	26	20	17	11	6	1	0	0
HER2 IHC 0, Chemo + pembro	82	70	58	50	37	25	17	13	7	5	4	3	1	0



No. at risk	0	2	4	6	8	10	12	14	16	18	20	22	24	26
HER2 Low, SG + pembro	133	122	106	84	60	46	32	22	17	11	5	5	2	0
HER2 Low, Chemo + pembro	136	118	98	71	50	33	23	16	14	8	3	1	0	0

PFS curves showed benefit with SG + pembro compared to chemo + pembro, regardless of HER2 expression status

BICR, blinded independent central review; chemo, chemotherapy; HER2, human epidermal growth factor receptor 2; HR, hazard ratio; IHC, immunohistochemistry; KM, Kaplan-Meier; mo, months; pembro, pembrolizumab; PFS, progression-free survival; SG, sacituzumab govitecan.

# Conclusions

- Consistent with the results of the primary analysis of ASCENT-04, median PFS was longer across all key participant subgroups with SG + pembro vs chemo + pembro<sup>1</sup>
  - All Trop-2 expression quartiles
  - Participants with both tBRCA WT and tBRCA mut status
  - HER2 IHC 0 and HER 2 low expression subgroups
- The results of these analyses are consistent with results from the ASCENT study, in which participants with metastatic TNBC treated with SG had longer PFS versus chemotherapy across all Trop-2, BRCA, and HER2 categories<sup>2,3</sup>
- Sample sizes in many subgroups were small and the analyses were descriptive, so caution should be used in interpreting these results

The results of this analysis reinforce the significant, clinically meaningful benefit of SG + pembro as a first-line treatment option for patients with previously untreated advanced PD-L1–positive TNBC across key Trop-2, tBRCA, and HER2 biomarker subgroups

**Chemo**, chemotherapy; **HER2**, human epidermal growth factor receptor 2; **mut**, mutant; **PD-L1**, programmed cell death ligand 1; **pembro**, pembrolizumab; **PFS**, progression-free survival; **SG**, sacituzumab govitecan; **tBRCA**, tumor BRCA; **TNBC**, triple-negative breast cancer; **WT**, wild-type.

1. Tolaney SM, et al. *N Engl J Med*. 2026;394:354-66. 2. Bardia A, et al. *J Clin Oncol*. 2024;42:1738-44. 3. Bardia A, et al. *Ann Oncol*. 2021;32:1148-56.

# Acknowledgments

- We thank the patients and their caregivers for their participation and commitment to clinical research
- Thank you to the clinical trial investigators and their devoted team members for contributing to the ASCENT-04 study
- This study was sponsored by Gilead Sciences, Inc.
- Medical writing support was provided by Ben Labbe, PhD, of Parexel, and funded by Gilead Sciences, Inc.
- Correspondence: Sara M Tolaney, Sara\_Tolaney@DFCI.HARVARD.EDU



A **plain language summary** of this presentation is available via QR code\* on this slide.

\*Copies of this slide deck obtained through Quick Response (QR) Code are for personal use only and may not be reproduced without permission from ASCO® or the author of these slides.