

# Epclusa<sup>®</sup> (SOF/VEL)

## Use in Renal Impairment

This document is in response to your request for information regarding the use of Epclusa<sup>®</sup> (sofosbuvir/velpatasvir [SOF/VEL]) in patients with renal impairment (eGFR <90 mL/min/1.73 m<sup>2</sup>), including those who require hemodialysis (HD).

Some data may be outside of the US FDA-approved prescribing information. In providing this data, Gilead Sciences, Inc. is not making any representation as to its clinical relevance or to the use of any Gilead product(s). For information about the approved conditions of use of any Gilead drug product, please consult the FDA-approved prescribing information.

**The full indication, important safety information, and boxed warnings are available at: [www.gilead.com/-/media/files/pdfs/medicines/liver-disease/epclusa/epclusa\\_pi](http://www.gilead.com/-/media/files/pdfs/medicines/liver-disease/epclusa/epclusa_pi).**

---

## Summary

### Product Labeling<sup>1</sup>

No dosage adjustment of SOF/VEL is recommended for patients with mild, moderate, or severe renal impairment, including ESRD requiring dialysis. No safety data are available in patients with both decompensated cirrhosis and severe renal impairment, including ESRD requiring dialysis. Additionally, no safety data are available in pediatric patients with renal impairment. Refer to RBV tablet prescribing information regarding use of RBV in patients with renal impairment.

### Clinical Studies of SOF/VEL Use in Renal Impairment

In a phase 2 study that evaluated participants with severe renal impairment, including those undergoing HD or PD, the SVR12 rate was 95%. AEs included headache, fatigue, nausea, vomiting, insomnia, and anemia.<sup>2</sup>

### Real-World Studies of SOF/VEL Use in Renal Impairment

A claims database study in patients with CKD found no difference in the risk of ESRD or HD between those treated with SOF-based (n=2042) vs non-SOF-based (n=431) regimens.<sup>3</sup>

In a prospective real-world study, 99.2% of patients (870/877) achieved SVR12 with SOF/VEL ± RBV, and rates were similar (99.1–100%) among subgroups according to baseline eGFR. Patients with baseline eGFR ≤60 mL/min/1.73 m<sup>2</sup> experienced significant increases in eGFR at EOT and EOF, whereas those with baseline eGFR >60 mL/min/1.73 m<sup>2</sup> experienced significant decreases in eGFR at both time points.<sup>4</sup>

A systematic review and meta-analysis of 21 studies evaluated SOF-based treatments in 717 patients with Stage 4 or 5 CKD. SVR12 was achieved by 97% of patients.<sup>5</sup>

In retrospective real-world studies conducted in patients with CKD, including those with severe renal impairment, approximately 95% to 100% of those treated with SOF/VEL ± RBV achieved SVR12.<sup>6,7</sup>

## Clinical Studies of SOF/VEL Use in Renal Impairment

### Study 4062: SOF/VEL in Participants Undergoing HD or PD<sup>2</sup>

#### Study design and demographics

Study 4062 was a phase 2, multicenter, open-label study that evaluated the efficacy and safety of SOF/VEL in 59 participants who were undergoing HD (n=54) or PD (n=5). Participants received SOF/VEL (400/100 mg) daily for 12 weeks, and SVR12 was assessed at Week 24. Pharmacokinetic levels were collected at each study visit while participants were receiving SOF/VEL. At baseline, the mean age was 60 years, 59% (35/59) were males, 29% (17/59) had compensated cirrhosis, and 22% (13/59) were treatment experienced. The mean duration of dialysis was 7.3 years, and 32% (19/59) had a history of kidney transplant.

#### Results

SVR12 was achieved in 95% of participants (56/59). Among the 3 participants who did not achieve SVR12, 1 participant with HCV GT 3 and cirrhosis relapsed, 1 discontinued the study due to low study drug adherence and subsequently relapsed, and 1 achieved SVR4 and died by suicide after the 12-week study period.

The most common ( $\geq 10\%$ ) AEs were headache (17%), fatigue (14%), nausea (14%), vomiting (14%), and insomnia (10%). Seven Grade 3 AEs and 11 SAEs occurred; none of these events were deemed related to study treatment. Grade 3 or 4 laboratory abnormalities (elevated Cr, n=15 [26%], elevated glucose, n=5 [8%], decreased Hgb, n=4 [7%], and elevated potassium, n=3 [5%]) were expected changes in participants undergoing dialysis, and elevated glucose values were seen in participants with diabetes. The area under the plasma concentration-time curve over the dosing interval for the predominant circulating metabolite of SOF (GS-331007) in participants undergoing HD or PD was 1719% (90% CI: 1617–1826%) higher than in participants with eGFR  $\geq 90$  mL/min who were enrolled in phase 2/3 studies.

---

## Real-World Studies of SOF/VEL Use in Renal Impairment

### US Claims Database Analysis<sup>3</sup>

#### Study design and demographics

The risk of ESRD or dialysis among patients with CKD and HCV treated with DAA regimens was evaluated using data sourced from a large US database (IQVIA PharMetrics Plus). The non-concurrent, prospective, observational cohort included claims for adult patients with DAA treatment for HCV infection, CKD diagnosis prior to HCV treatment initiation, and  $\geq 12$  months of enrollment prior to HCV treatment start that occurred between January 1, 2006, and March 30, 2019. There was a higher proportion of patients with severe CKD in the non-SOF-based treatment cohort (eg, OBV/PTV/RTV + DSV, OBV/PTV/RTV, EBR/GZR, and GLE/PIB) than in the SOF-based treatment cohort (eg, SOF, LDV/SOF, and SOF/VEL).

**Table 1. Baseline Demographics and Disease Characteristics (Telep et al)<sup>3</sup>**

Key Demographics and Characteristics	SOF-Based (n=2042)	Non-SOF-Based (n=431)
Age, 18–34/35–44/45–54/ 55–64/65–74/≥75 years, n (%)	27 (1)/59 (3)/295 (14)/ 1300 (64)/331 (16)/30 (1)	14 (3)/11 (3)/64 (15)/ 241 (56)/89 (21)/12 (3)
Male, n (%)	1424 (70)	293 (68)
CKD stage, mild/moderate/severe/unspecified, n (%)	347 (17)/696 (34)/ 123 (6)/876 (43)	65 (15)/161 (37)/ 68 (16)/137 (32)

## Efficacy

The crude incidence rates of ESRD or dialysis were highest for patients with severe CKD and in the cohort that did not receive SOF. After adjustment for age, sex, and significant covariates and PS-weighting by treatment, there was no difference in the risk of ESRD or dialysis between SOF- and non-SOF-based regimens. Similar results were observed when stratified by CKD stage.

**Table 2. ESRD/Dialysis Rates in SOF-Based and Non-SOF-Based DAA Cohorts, Stratified by Baseline CKD Stage<sup>a</sup> (Telep et al)<sup>3</sup>**

	Moderate CKD		Severe CKD	
	SOF-Based (n=696)	Non-SOF-Based (n=161)	SOF-Based (n=123)	Non-SOF-Based (n=68)
Number of events	56	8	25	18
Time at risk, PY	1072	138	165	55
Unadjusted rate of ESRD/ dialysis per 100 PY (95% CI)	5.22 (3.95–6.78)	5.78 (2.49–11.39)	15.11 (9.78–22.31)	32.66 (19.35–51.61)
Adjusted and PS-weighted, <sup>b</sup> HR (95% CI)	0.91 (0.41–2.01)	1	0.83 (0.35–2.02)	1

Abbreviation: PY=person-years; HR=hazard ratio

<sup>a</sup>Results were stratified by participants' baseline CKD stage, where counts of events for each stratum (SOF-based vs non-SOF-based regimens) were ≥5.

<sup>b</sup>HR adjusted for age group, sex, prior diagnosis for acute kidney injury, anemia, hepatic encephalopathy, coronary atherosclerosis, arrhythmia, liver necrosis, essential hypertension, and prior prescriptions for calcium channel blockers or diabetes medication; weighting was based on treatment PS.

## Taiwan Prospective Cohort Study of SOF/VEL ± RBV<sup>4</sup>

### Study design and demographics

A single-center, prospective cohort study was conducted in Taiwan to evaluate the changes in eGFR in 953 participants with chronic HCV. Participants with detectable HCV RNA (LLoD=15 IU/mL) who received SOF/VEL (400/100 mg) ± RBV for 12 weeks were included in the study. Participants on HD or PD were excluded. The eGFR was assessed at baseline, EOT, and EOF (defined as 12 weeks after completion of treatment). SVR12 rates were assessed in both the EP (defined as participants who received ≥1 dose of SOF/VEL ± RBV) and PP population (defined as participants who received ≥1 dose of SOF/VEL ± RBV and had HCV RNA data at Week 12 after treatment). Among the total participants, mean eGFR was 85.57 mL/min/1.73 m<sup>2</sup>; a total of 130 patients had eGFR ≤60 mL/min/1.73 m<sup>2</sup> and 823 had eGFR >60 mL/min/1.73 m<sup>2</sup> at baseline (Table 3).

**Table 3. Baseline Demographics and Disease Characteristics (Su et al)<sup>4</sup>**

Key Demographics and Characteristics	SOF/VEL ± RBV		P-Value
	Baseline eGFR ≤60 mL/min/1.73 m <sup>2</sup> (n=130)	Baseline eGFR >60 mL/min/1.73 m <sup>2</sup> (n=823)	
Age, mean (SD), years	74.24±9.99	62.69±14.31	<0.001
Male, n	65	416	0.98
RBV used, n	12	27	<0.05
FIB-4, ≤3.25/>3.25, n	85/45	641/182	<0.05
HCV RNA, <800,000/≥800,000 IU/mL, n	55/75	268/555	<0.05
Albumin, mean (SD), g/dL	4±0.54	4.21±0.41	<0.001
Total bilirubin, mean (SD), mg/dL	0.88±0.56	0.95±0.75	0.23
AST, mean (SD), U/L	49.88±39.14	52.44±46.99	0.5
ALT, mean (SD), U/L	48.81±42.07	64.03±69.68	<0.001
Cr, mean (SD), mg/dL	1.43±0.45	0.83±0.18	<0.001
eGFR, mean (SD), mL/min/1.73 m <sup>2</sup>	47.89±10.25	91.52±22.06	<0.001

## Effectiveness

Overall, SVR12 was achieved in 99.2% of participants (870/877) in the PP analysis and in 91.3% (870/953) in the EP analysis; 4 participants experienced virological relapse, and 3 participants did not respond. The SVR12 rates according to baseline eGFR were also reported (Table 4). SVR12 rates did not differ significantly between participants with eGFR ≤60 mL/min/1.73 m<sup>2</sup> and those with eGFR >60 mL/min/1.73 m<sup>2</sup> in subgroups according to age, sex, HBV co-infection, history of hepatocellular carcinoma, prior treatment experience, history of diabetes mellitus, baseline HCV RNA level, and baseline FIB-4 score.


**Table 4. SVR12 Rates According to Baseline eGFR (Su et al)<sup>4</sup>**

HCV RNA < LLoD, n/N (%)		SOF/VEL ± RBV		
		Baseline eGFR <30 mL/min/1.73 m <sup>2</sup> (n=9)	Baseline eGFR 30–60 mL/min/1.73 m <sup>2</sup> (n=121)	Baseline eGFR >60 mL/min/1.73 m <sup>2</sup> (n=823)
After treatment	SVR12 (EP)	8/9 (88.9)	110/121 (90.9)	752/823 (91.4)
	SVR12 (PP)	8/8 (100)	110/110 (100)	752/759 (99.1)

## Safety

Participants with baseline eGFR ≤60 mL/min/1.73 m<sup>2</sup> showed a significant increase from baseline in eGFR at EOT ( $P<0.001$ ) and EOF ( $P<0.05$ ). Participants with baseline eGFR >60 mL/min/1.73 m<sup>2</sup> experienced significant decreases from baseline in eGFR at EOT ( $P<0.05$ ) and EOF ( $P<0.001$ ; Figure 1). In multivariable analyses, participants with higher albumin levels were less likely to experience reductions of >5% in eGFR at EOT (OR, 0.45; 95% CI: 0.28–0.69;  $P<0.001$ ), whereas participants with a baseline eGFR >60 mL/min/1.73 m<sup>2</sup> were more likely to experience eGFR reductions at EOF than those with a baseline eGFR ≤60 mL/min/1.73 m<sup>2</sup> (OR, 1.58; 95% CI: 1.03–2.45;  $P<0.05$ ).

**Figure 1. Change in eGFR (Su et al)<sup>4</sup>**



<sup>a</sup>P value for comparison with baseline eGFR.

<sup>b</sup>P value for comparison between eGFR at EOT and eGFR at EOF.

Note: In the baseline eGFR >60 mL/min/1.73 m<sup>2</sup> cohort, n=823 at baseline, n=652 at EOT, and n=762 at EOF. In the baseline eGFR ≤60 mL/min/1.73 m<sup>2</sup> cohort, n=130 at baseline, n=107 at EOT, and n=119 at EOF.

No SOF/VEL treatment-related deaths occurred during the study. AEs reported in >3% of participants and laboratory AEs are summarized in Table 5.

**Table 5. Summary of Safety Results (Su et al)<sup>4</sup>**

Events, n (%)		SOF/VEL ± RBV		
		Baseline eGFR ≤60 mL/min/1.73 m <sup>2</sup> (n=130)	Baseline eGFR >60 mL/min/1.73 m <sup>2</sup> (n=823)	Total (N=953)
AEs >3%	Fatigue	9 (6.9)	36 (4.4)	45 (4.7)
	Abdominal discomfort	5 (3.8)	38 (4.6)	43 (4.5)
	Skin itching	5 (3.8)	30 (3.6)	35 (3.7)
Increased total bilirubin	1.5–3 × ULN	6 (4.6)	18 (2.2)	24 (2.5)
	>3 × ULN	0	6 (0.7)	6 (0.6)
Increased AST	3–5 × ULN	2 (1.5)	10 (1.2)	12 (1.3)
	>5 × ULN	1 (0.8)	1 (0.1)	2 (0.2)
Increased ALT	3–5 × ULN	2 (1.5)	4 (0.5)	6 (0.6)
	>5 × ULN	0	3 (0.4)	3 (0.3)

Abbreviation: ULN=upper limit of normal.

## Systematic Review and Meta-Analysis<sup>5</sup>

### Study design and demographics

A systematic review and meta-analysis of 21 studies included 717 patients with HCV and Stage 4 or 5 CKD who were treated with SOF-based regimens. Most studies included patients with a history of cirrhosis, and 4 studies included those with decompensated cirrhosis. The following SOF-based regimens were included: SOF + SMV ± RBV, SOF + PEG-IFN, SOF + RBV, SOF + DCV ± RBV, and LDV/SOF ± RBV. The dosing of SOF varied by study: SOF 400 mg once daily, 200 mg once daily, 400 mg every other day, and 400 mg three times weekly. Treatment was administered for 8, 12, 16, or 24 weeks (where reported). Included patients had a mean/median age of 35 to 62 years, and 58% were receiving HD or PD; in 18 studies, the percentages of patients with HCV GTs 1, 2, and 3 were 67%, 8%, and 20%, respectively.

## Efficacy

SVR12/24 rates were significantly higher among the 10 studies that did not utilize RBV in addition to an SOF-based regimen than among those that included RBV in the treatment regimen (99% vs 94%;  $P=0.035$ ). Seven studies included data that allowed comparisons of efficacy outcomes between patients with and those without cirrhosis. Patients without cirrhosis tended to have higher SVR12/24 rates than those with cirrhosis, although the difference was not significant (relative risk, 0.93; 95% CI: 0.85–1.02;  $I^2=33\%$ ).

## Safety

The pooled SAE rate for the 16 studies with available data was 4.8% (95% CI: 2.1–10.3%;  $I^2=60\%$ ). There was no significant difference in the incidence of SAEs among patients who received full-dose and decreased-dose SOF (8.8% vs 2.9%;  $P=0.13$ ). Of the four studies that reported changes in kidney function, eGFR values were generally unchanged during treatment, and 2 patients discontinued HD as a result of eGFR improvement.

## Taiwan Retrospective Cohort Study of SOF/VEL ± RBV<sup>6</sup>

### Study design and demographics

A multicenter, retrospective study was conducted to assess the effectiveness and tolerance of SOF/VEL ± RBV in patients infected with chronic HCV and CKD Stage 4 (eGFR <30 mL/min/1.73 m<sup>2</sup>) or Stage 5 (eGFR <15 mL/min/1.73 m<sup>2</sup>) with decompensated or compensated liver disease. Data were collected from fifteen academic centers in Taiwan between July 2019 and March 2020. Patients aged ≥20 years with chronic HCV infection (detectable HCV antibody and quantifiable serum HCV RNA for ≥6 months) were included in this analysis. The exclusion criteria included off-label use of SOF/VEL, solid organ transplant (other than failed renal transplant), and virological failures with NS5A-containing DAA regimens.

**Table 6. Baseline Demographics and Disease Characteristics (Liu et al)<sup>6</sup>**

Key Demographics and Characteristics		SOF/VEL (n=181)	SOF/VEL + RBV (n=10)	Total (N=191)
eGFR <sub>CKD-EPI</sub> , median (range), mL/min/1.73 m <sup>2</sup>		8 (2–29)	24 (5–29)	8 (2–29)
CKD stage, Stage 4/Stage 5/HD or PD, n (%)		56 (30.9)/125 (69.1)/ 111 (97.4)/3 (2.6)	7 (70)/ 3 (30)/ 3 (100)/0	63 (33)/128 (67)/ 114 (97.4)/3 (2.6)
History of renal transplantation, n (%)		8 (4.4)	0	8 (4.2)
Fibrosis stage, n (%)	F0–1/F2/F3/F4 <sup>a</sup>	52 (28.7)/92 (50.8)/ 20 (11)/17 (9.4)	0/0/0/10 (100)	52 (27.2)/92 (48.2)/ 20 (10.5)/27 (14.1)
	Compensated/decompensated	17 (100)/0	0/10 (100)	17 (63)/10 (37)

Abbreviation: CKD-EPI=Chronic Kidney Disease Epidemiology Collaboration equation.

<sup>a</sup>F4 was defined as a fibrosis index-based score >3.25 and an AST-to-platelet ratio index score >2.

## Effectiveness

After 12 weeks of treatment, all patients with available data (n=187) had HCV RNA levels <LLOQ. SVR12 rates are shown in Table 7. Among patients with compensated liver disease, 94.6% of patients (53/56) with CKD Stage 4 and 95.2% (119/125) with Stage 5 achieved SVR12. Among patients with decompensated liver disease, 100% of patients (7/7) with CKD Stage 4 and 66.7% (2/3) with Stage 5 achieved SVR12.



**Table 7. Summary of Patients Who Achieved SVR12 (Liu et al)<sup>6</sup>**

Outcomes, n/N (%)	SOF/VEL (n=181)	SOF/VEL + RBV (n=10)	Total (N=191)
HCV RNA <LLOQ (15 IU/mL) at Week 12 of treatment <sup>a</sup>	178/178 (100)	9/9 (100)	187/187 (100)
SVR12 (EP <sup>b</sup> )	172/181 (95)	9/10 (90)	181/191 (94.8)
SVR12 (PP <sup>c</sup> )	172/172 (100)	9/9 (100)	181/181 (100)

<sup>a</sup>Data were not available from 4 patients who died or discontinued treatment prematurely.

<sup>b</sup>The EP was defined as patients who received ≥1 dose of SOF/VEL ± RBV.

<sup>c</sup>The PP population was defined as patients with available SVR12 data.

## Safety

Treatment discontinuation due to general weakness was reported by 1 patient with Stage 4 CKD and Stage F2 fibrosis (Table 8). Changes in eGFR levels in patients not on maintenance dialysis are summarized in Table 9.

**Table 8. Summary of Safety Results (Liu et al)<sup>6</sup>**

Safety Outcomes, n (%)	SOF/VEL (n=181)	SOF/VEL + RBV (n=10)	Total (N=191)
Any AE	110 (60.8)	9 (90)	119 (62.3)
SAE	18 (9.9)	2 (20)	20 (10.5)
Treatment-related SAE	0	0	0
SOF/VEL discontinuation due to TEAE	1 (0.6)	0	1 (0.5)
Death <sup>a</sup>	6 (3.3)	1 (10)	7 (3.7)

Abbreviation: TEAE=treatment-emergent adverse event.

<sup>a</sup>No deaths were considered to be related to SOF/VEL or RBV.

**Table 9. Changes in eGFR Levels in Patients Not on Maintenance Dialysis (Liu et al)<sup>6</sup>**

eGFR	CKD Stage 4 (n=56)	CKD Stage 5 (n=11)
At baseline, mean (SD), mL/min/1.73 m <sup>2</sup>	25.5 (3.3)	7.3 (3.3)
At 12 weeks after treatment, mean (SD), mL/min/1.73 m <sup>2</sup>	27 (6.1)	7 (3.3)
P-value for baseline vs 12 weeks after treatment	0.06	0.65

## Chinese Retrospective Study in Patients With CKD<sup>7</sup>

### Study design and demographics

A retrospective study was conducted to assess the effectiveness and safety of SOF/VEL ± RBV in 75 patients with chronic HCV and CKD (eGFR <90 mL/min/1.73 m<sup>2</sup>). Data were collected from a single center in China between June 2018 and May 2022. Patients who received SOF/VEL ± RBV (with the RBV dose varying according to weight and renal function) for 12 weeks were included in the study. The primary endpoint was change in renal function over 12 weeks. SVR12 rates and AEs were also evaluated.

**Table 10. Baseline Demographics and Disease Characteristics (Yang et al)<sup>7</sup>**

Key Demographics and Characteristics	SOF/VEL ± RBV (N=75)
Age, mean (range), years	52 (30–85)
Male, n (%)	58 (77.3)
GT 1b/2a/3a/3b/6n/unknown, n	1/4/26/37/4/3
CKD Stage 1/2/3a/3b/4/5, n (%)	0/51 (68)/11 (14.6)/1 (1.3)/4 (5.3)/8 (10.7)

Key Demographics and Characteristics		SOF/VEL ± RBV (N=75)
Cirrhosis status, n (%)	Compensated	15 (20)
	Decompensated	36 (48)

## Effectiveness

Overall, SVR12 (HCV RNA <20 IU/mL) was achieved in 98.7% of patients (74/75). The 1 patient who did not achieve SVR12 had decompensated cirrhosis and relapsed liver cancer.

## Safety

A worsening in renal function to a higher stage of CKD occurred in 8% of patients (6/75) (Table 11). Changes in laboratory parameters (eGFR, ALT, and AST) were observed from baseline to off-treatment (Table 12).

**Table 11. Change in CKD Stages From Baseline to Off-Treatment (Yang et al)<sup>2</sup>**

Baseline CKD Stage	Off-Treatment CKD Stage, n (%)					
	Stage 1	Stage 2	Stage 3a	Stage 3b	Stage 4	Stage 5
Stage 2 (n=51)	14 (27.5)	34 (66.7)	2 (3.9)	1 (2)	0	0
Stage 3a (n=11)	1 (9.1)	4 (36.4)	4 (36.4)	2 (18.2)	0	0
Stage 3b (n=1)	0	1 (100)	0	0	0	0
Stage 4 (n=4)	1 (25)	0	0	1 (25)	1 (25)	1 (25)
Stage 5 (n=8)	0	0	0	0	0	8 (100)
<div>Dark purple cells=Improved</div> <div>White cells=No change</div> <div>Gray cells=Worsened</div>						

**Table 12. Changes in Laboratory Parameters (Yang et al)<sup>2</sup>**

Laboratory Parameters	SOF/VEL ± RBV (N=75)		
	Baseline	Off-Treatment	P-Value
eGFR, mean ± SD, mL/min	63.19±3.01	72.71±4.73	0.078
ALT, mean ± SD, U/L	51.36±35.078	25.3±35.641	<0.05
AST, mean ± SD, U/L	58.1±35.686	42.7±75.758	<0.05

Twenty-two patients (29.3%) reported AEs during treatment with SOF/VEL ± RBV, including hemolysis (n=16), dizziness (n=2), fatigue (n=2), pruritus (n=1), and rash (n=1). All AEs were resolved following supportive treatment, and no SAEs were reported.

## References

1. Enclosed. Gilead Sciences Inc, EPCLUSA® (sofosbuvir and velpatasvir) tablets, for oral use. US Prescribing Information. Foster City, CA
2. Borgia SM, Dearden J, Yoshida EM, et al. Sofosbuvir/velpatasvir for 12 weeks in hepatitis C virus-infected patients with end-stage renal disease undergoing dialysis. *J Hepatol*. 2019;71(4):660-665. <https://www.ncbi.nlm.nih.gov/pubmed/31195062>



3. Telep LE, Sise M, Force L, Gaggar A, Chokkalingam AP. Risk of end-stage renal disease among hepatitis C patients with chronic kidney disease treated with and without sofosbuvir [Poster FRI-441]. Paper presented at: European Association for the Study of the Liver (EASL): The Digital International Liver Congress 27-29 August, 2020.
4. Su P, Chang T, Tung S, et al. Changes in renal function in patients with chronic hepatitis C treated with sofosbuvir-velpatasvir. *Advances in Digestive Medicine*. 2023;10(3):150-158.
5. Li M, Chen J, Fang Z, Li Y, Lin Q. Sofosbuvir-Based Regimen Is Safe and Effective for Hepatitis C Infected Patients with Stage 4-5 Chronic Kidney Disease: A Systematic Review and Meta-Analysis. *Virology*. 2019;16:34. <https://www.ncbi.nlm.nih.gov/pubmed/30871566>
6. Liu CH, Chen CY, Su WW, et al. Sofosbuvir/velpatasvir with or without low-dose ribavirin for patients with chronic hepatitis C virus infection and severe renal impairment. *Gut*. 2021.
7. Yang Y, Gao J, bai J, et al. Sofosbuvir/Velpatasvir with or without Ribavirin for HCV infected patients who had chronic kidney disease —A real world experience from Southwest China. Paper presented at: Asian Pacific Association for the Study of the Liver (APASL); February 15 – 19, 2023; Taipei, Taiwan.

---

## Abbreviations

AE=adverse event  
CKD=chronic kidney disease  
DAA=direct-acting antiviral  
DCV=daclatasvir  
DSV=dasabuvir  
EBR=elbasvir  
EOF=end of follow-up  
EOT=end of treatment  
EP=evaluable population  
ESRD=end-stage renal disease  
FIB-4=fibrosis-4 index  
GLE=glecaprevir

GS-331007=predominant circulating metabolite of SOF  
GT=genotype  
GZR=grazoprevir  
HD=hemodialysis  
LDV=ledipasvir  
LLOD=lower limit of detection  
LLOQ=lower limit of quantification  
OBV=ombitasvir  
OR=odds ratio  
PD=peritoneal dialysis  
PEG-IFN=pegylated interferon

PIB=pibrentasvir  
PP=per-protocol  
PS=propensity score  
PTV=paritaprevir  
RBV=ribavirin  
RTV=ritonavir  
SAE=serious adverse event  
SMV=simeprevir  
SOF=sofosbuvir  
SVR4/12/24=sustained virologic response 4/12/24 weeks after end of treatment  
VEL=velpatasvir

---

## Product Label

For the full indication, important safety information, and boxed warning(s), please refer to the Epclusa US Prescribing Information available at:

[www.gilead.com/-/media/files/pdfs/medicines/liver-disease/epclusa/epclusa\\_pi](http://www.gilead.com/-/media/files/pdfs/medicines/liver-disease/epclusa/epclusa_pi).

## Follow-Up

For any additional questions, please contact Gilead Medical Information at:

☎ 1-866-MEDI-GSI (1-866-633-4474) or 🌐 [www.askgileadmedical.com](http://www.askgileadmedical.com)

## Adverse Event Reporting

Please report all adverse events to:

Gilead Global Patient Safety ☎ 1-800-445-3235, option 3 or

🌐 [www.gilead.com/utility/contact/report-an-adverse-event](http://www.gilead.com/utility/contact/report-an-adverse-event)

FDA MedWatch Program by ☎ 1-800-FDA-1088 or ✉ MedWatch, FDA, 5600 Fishers Ln, Rockville, MD 20852 or 🌐 [www.accessdata.fda.gov/scripts/medwatch](http://www.accessdata.fda.gov/scripts/medwatch)

## Data Privacy

The Medical Information service at Gilead Sciences may collect, store, and use your personal information to provide a response to your medical request. We may share your information with other Gilead Sciences colleagues to ensure that your request is addressed appropriately. If you report an adverse event or concern about the quality of a Gilead or Kite product, we will need to use the information you have given us in order to meet our regulatory requirements in relation to the safety of our medicines.

It may be necessary for us to share your information with Gilead's affiliates, business partners, service providers, and regulatory authorities located in countries besides your own. Gilead Sciences has implemented measures to protect the personal information you provide. Please see the Gilead Privacy Statement ([www.gilead.com/privacy-statements](http://www.gilead.com/privacy-statements)) for more information about how Gilead handles your personal information and your rights. If you have any further questions about the use of your personal information, please contact [privacy@gilead.com](mailto:privacy@gilead.com).

EPCLUSA, GILEAD, and the GILEAD logo are registered trademarks of Gilead Sciences, Inc., or its related companies.

© 2025 Gilead Sciences, Inc.